

Claims

1. A composition comprising:

(a) an oxygen labile active agent, and

(b) a plant extract selected from the group
consisting of Chrysanthellum extract, Neem extract,
Lanatellys extract, Bacopa Monnieri extract, and Olive
leaf extract.

2. A composition of claim 1, wherein said plant
extract is Chrysanthellum extract.

3. A composition of claim 1, wherein said plant
extract is Neem extract.

4. A composition of claim 1, wherein said plant
extract is Lanatellys extract.

5. A composition of claim 1, wherein said plant
extract is Bacopa Monnieri extract.

6. A composition of claim 1, wherein said plant
extract is Olive leaf extract.

7. A composition of claim 1, wherein said oxygen
labile active agent is selected from the group consisting
of ascorbic acid, tocotrienol, hydroquinone, ubiquinone,
and dihydrolipoic acid.

8. A composition of claim 1, wherein said oxygen
labile active agent is a retinoid.

9. A composition of claim 1, wherein said oxygen
labile active agent is retinol.

10. A composition of claim 9, wherein said composition comprises (i) from about 0.01% to about 1%, by weight, of said retinol and (ii) from about 0.1% to about 10%, by weight, of said plant extract.

5

11. A composition comprising:
(a) an oxygen labile active agent,
(b) an isoascorbic acid derivative,
(c) a tocopherol derivative, and
10 (d) a plant extract selected from the group consisting of Chrysanthellum extract, Neem extract, Lanatellys extract, Bacopa Monnieri extract, and Olive leaf extract.

15

12. A composition of claim 1, wherein said plant extract is Chrysanthellum extract.

13. A composition of claim 1, wherein said plant extract is Neem extract.

20

14. A composition of claim 1, wherein said plant extract is Lanatellys extract.

15. A composition of claim 1, wherein said plant extract is Bacopa Monnieri extract.

25

16. A composition of claim 1, wherein said plant extract is Olive leaf extract.

30

17. A composition of claim 1, wherein said composition comprises:

(a) about 0.001 to about 20%, by weight, of said oxygen labile active agent,

(b) about 0.001% to about 0.5%, by weight, of said isoascorbic acid derivative, and

35

(c) about 0.1% to about 1%, by weight, of said tocopherol derivative; and

(d) about 0.1% to about 10%, by weight, of said plant extract.

5

18. A composition of claim 1, wherein said oxygen labile active agent is selected from the group consisting of ascorbic acid, tocotrienol, hydroquinone, ubiquinone, and dihydrolipoic acid.

10

19. A composition of claim 1, wherein said oxygen labile active agent is a retinoid.

20. A composition of claim 1, wherein said oxygen labile active agent is retinol.

15